

EXTENDED CANONICAL ALGEBRAS AND FUCHSIAN SINGULARITIES

HELMUT LENZING AND JOSÉ A. DE LA PEÑA

1. Introduction

1.1. Main results. Let k be any field. Consider the hereditary algebra $A = k\Delta$ associated to a finite connected quiver without oriented cycles. A fundamental fact in Representation Theory of Algebras is the distinction of the representation type of A : A is *representation-finite* (that is, there are only finitely many indecomposable A -modules, up to isomorphism) exactly when the underlying graph $|\Delta|$ of Δ is of Dynkin type; A is *tame* (that is, for each $d \in \mathbb{N}$, the indecomposable d -dimensional A -modules may be classified in a finite number of one-parameter families of modules) exactly when $|\Delta|$ is of extended Dynkin type; in the remaining cases A is *wild* (that is, there is a embedding $\text{mod-}k\langle x, y \rangle \rightarrow \text{mod-}A$ which preserves indecomposability and isomorphism classes from the category of finite dimensional modules over the ring in two non-commuting indeterminates into the category of finite dimensional A -modules). Consider the Auslander-Reiten translation τ_A in $\text{mod-}A$ and P an indecomposable projective A -module. In case A is tame representation-infinite, the sequence of modules $(\tau_A^{-n}P)_n$ is well defined and the algebra $R(A, P) = \bigoplus_{n=0}^{\infty} \text{Hom}_A(P, \tau_A^{-n}P)$ is an infinite dimensional positively $\mathbb{Z}$ -graded surface singularity. The algebra $R(A, P)$ reflects many properties of A and P and has a particularly interesting structure, as shown in [12]:

Assume $A = \mathbb{C}\tilde{\Delta}$ is a tame hereditary algebra where $\tilde{\Delta}$ extends the Dynkin type $|\Delta|$. Let P the indecomposable projective associated to the vertex in $\tilde{\Delta} \setminus \Delta$. Then $R(A, P)$ is isomorphic to the algebra of invariants $\mathbb{C}[x, y]^G$, where $G \subset SL(2, \mathbb{C})$ is a binary polyhedral group of type $|\Delta|$. Accordingly the completion of the graded algebra $R(A, P)$ is isomorphic to the surface singularity of type $|\Delta|$.

In [34], Ringel introduced the canonical algebras $C = C(p, \lambda)$ depending on a weight sequence $p = (p_1, \dots, p_t)$ of positive integers and a parameter sequence $\lambda =$

$(\lambda_3, \dots, \lambda_t)$ of pairwise distinct non-zero elements from k . In [11] it was shown that $\text{mod-}C$ is derived equivalent to $\text{coh}(\mathbb{X})$ the category of coherent sheaves on a weighted projective line $\mathbb{X} = \mathbb{X}(p, \lambda)$.

One of the aims of this work is the introduction of a class of algebras with related interesting properties. Let P be an indecomposable projective module over a canonical algebra $C = C(p, \lambda)$, the one-point extension $A = C[P]$ defined as the matrix algebra

$$\begin{bmatrix} k & 0 \\ P & C \end{bmatrix}$$

is called an *extended canonical algebra*. In section 2 we show that for two indecomposable projective C -modules P and P' , the algebras $C[P]$ and $C[P']$ are derived equivalent. Moreover, if C is of tame type, the extended canonical algebra is derived equivalent to a wild hereditary or a wild canonical algebra, so essentially $C[P]$ belongs to a well-studied class of algebras. There are interesting phenomena arising when C (and hence A) is of wild type.

Consider the *Coxeter transformation* of A as an automorphism $\varphi_A: K_0(A) \rightarrow K_0(A)$ of the Grothendieck group of A , given on the classes of indecomposable projective modules by the formula $\varphi_A([P(S)]) = -[I(S)]$, where $P(S)$ (resp. $I(S)$) is the projective cover (resp. injective envelope) of a simple module S . The characteristic polynomial $f_A(T)$ of φ_A is called the *Coxeter polynomial* of A .

Let $\chi_{\mathbb{X}} = 2 - \sum_{i=1}^t (1 - 1/p_i)$ be the (orbifold) Euler characteristic of $\mathbb{X}$. As shown in [12], for $\chi_{\mathbb{X}} > 0$ the classification problem of $\text{coh}(\mathbb{X})$ is related to the problem of classifying the Cohen-Macaulay modules over a simple surface singularity and is in fact equivalent to the problem of classifying the graded Cohen-Macaulay modules over a corresponding quasi-homogeneous singularity. Assume $\chi_{\mathbb{X}} < 0$. By [23, 25] we know that for $A = C[P]$ an extended canonical algebra

$$f_A(T) = P_C(T)f_C(T)$$

where $P_C(T)$ is the Hilbert-Poincaré series of the positively graded algebra $R(p, \lambda) = \bigoplus_{n=0}^{\infty} \text{Hom}_C(M, \tau_C^n M)$, where M is a rank one not preprojective C -module. Equivalently, $R(p, \lambda) = \bigoplus_{n=0}^{\infty} \text{Hom}(\mathcal{O}, \tau_{\mathbb{X}}^n \mathcal{O})$, where $\mathcal{O}$ is the structure sheaf on $\mathbb{X}$. Recall from [21, 23] that in case $k = \mathbb{C}$, we can interpret $R(p, \lambda)$ as an algebra of entire automorphic forms associated to the action of a suitable *Fuchsian group* of the first kind, acting on the upper half plane $\mathbb{H}_+$.

From [21], we know that the k -algebra $R = R(p, \lambda)$ is commutative, graded integral Gorenstein, in particular Cohen-Macaulay, of Krull dimension two. The complexity of the surface singularity R is described by the triangulated category

$$D_{\text{Sg}}^{\mathbb{Z}}(R) = \frac{D^b(\text{mod}^{\mathbb{Z}}\text{-}R)}{D^b(\text{proj}^{\mathbb{Z}}\text{-}R)},$$

where $\text{mod}^{\mathbb{Z}}\text{-}R$ (resp. $\text{proj}^{\mathbb{Z}}\text{-}R$) denotes the category of finitely generated (resp. finitely generated projective) $\mathbb{Z}$ -graded R -modules. This category was considered by Buchweitz [5] and Orlov [30], see also Krause's account [19] for a related, but slightly different approach. For $\chi_{\mathbb{X}} > 0$, where the weight type of $\mathbb{X}$ determines a Dynkin quiver Δ , Kajiura, Saito, Takahashi and Ueda [17] have shown that $D_{\text{Sg}}^{\mathbb{Z}}(R)$ is equivalent to the derived category of finite dimensional modules over the path algebra $k\Delta$. For $\chi_{\mathbb{X}} = 0$, the algebra R hence $D_{\text{Sg}}^{\mathbb{Z}}(R)$ is not defined, but a close variant $D_{\text{Sg}}^{\mathbb{L}(p)}(S)$, as shown by Ueda [35], is equivalent to the derived category $D^b(\text{coh}(\mathbb{X}))$ of coherent sheaves on a weighted projective line which is tubular, that is, has weight type $(2, 3, 6)$, $(2, 4, 4)$, $(3, 3, 3)$ or $(2, 2, 2, 2)$.

In section 3 we deal with the case $\chi_{\mathbb{X}} < 0$ and prove that this category, as first observed by Saito and Takahashi (for the field $\mathbb{C}$ of complex numbers), is described in the following way.

Theorem 1. *Let k be an algebraically closed field. Assume $\chi_{\mathbb{X}} < 0$ and let R be the positively $\mathbb{Z}$ -graded surface singularity attached to $\mathbb{X}$. Then there exists a tilting object $\bar{T}$ in the triangulated category $D_{\text{Sg}}^{\mathbb{Z}}(R)$ whose endomorphism ring is isomorphic to an extended canonical algebra $C[P]$, where C is the canonical algebra associated with $\mathbb{X}$.*

It follows that the categories $D_{\text{Sg}}^{\mathbb{Z}}(R) = \mathcal{T}$ and $D^b(\text{mod}(C[P]))$ are equivalent as triangulated categories. In section 3.9 we further introduce the concept of Coxeter-Dynkin algebras and establish their relationship to the Coxeter-Dynkin diagrams from singularity theory.

More precise information on the structure of the ring $R(p, \lambda)$ is obtained by a closer examination of the spectral properties of the Coxeter transformation of the extended canonical algebra $C[P]$.

A sequence of weights $p = (p_1, \dots, p_t)$ will always satisfy $p_1 \leq p_2 \leq \dots \leq p_t$. We consider the lexicographical ordering of sequences $(p_1, \dots, p_t) \leq (q_1, \dots, q_s)$ if $t = s$

and $p_i \leq q_i$ for $1 \leq i \leq t$. Extend the relation $p \leq q$ (and say that q *dominates* p) to weight sequences of (possibly) different length by adding 1's if necessary. The following result is shown in section 4 based on techniques developed in [24] and will be fundamental in the proof of the main results.

Theorem 2. *Let $A = C[P]$ be an extended canonical algebra of the wild canonical algebra $C = C(p, \lambda)$. The following happens:*

- (a) $f_A(T)$ has at most 4 roots not in $\mathbb{S}^1$.
- (b) The roots of $f_A(T)$ lie on the unit circle $\mathbb{S}^1$ if and only if the weight sequence p belongs to the 38-member list determined by all $p < q$ with q belonging to the following critical list:
 - ($t = 3$): $(2, 3, 11), (2, 4, 9), (2, 5, 8), (2, 6, 7),$
 $(3, 3, 8), (3, 4, 7), (3, 5, 6),$
 $(4, 4, 6), (4, 5, 5),$
 - ($t = 4$): $(2, 2, 2, 7), (2, 2, 3, 6), (2, 3, 4, 4), (3, 3, 3, 4),$
 - ($t = 5$): $(2, 2, 2, 2, 5), (2, 2, 2, 3, 4), (2, 2, 3, 3, 3),$
 - ($t = 6$): $(2, 2, 2, 2, 2, 3),$
 - ($t = 7$): $(2, 2, 2, 2, 2, 2, 2).$

We shall say that the algebra $R(p, \lambda)$ (and also the weight sequence p) is *formally n -generated* if

$$P_C(T) = \frac{\prod_{i=1}^{n-2} (1 - T^{c_i})}{\prod_{j=1}^n (1 - T^{d_j})}$$

for certain natural numbers $c_1, \dots, c_{n-2}$ and $d_1, \dots, d_n$, all ≥ 2 . The algebra $R(p, \lambda)$ (and also the weight sequence p) is *formally a complete intersection* if $P_C(T)$ is a rational function $f_1(T)/f_2(T)$, where each $f_i(T)$ is a product of cyclotomic polynomials.

Theorem 3. *Let $C = C(p, \lambda)$ be a wild canonical algebra with weight sequence $(p_1, \dots, p_t)$ and $A = C[P]$ be an extended canonical algebra. The following are equivalent:*

- (a) $R(p, \lambda)$ is formally 3- or 4-generated
- (b) $R(p, \lambda)$ is formally a complete intersection

(c) *The roots of $f_A(T)$ lie on $\mathbb{S}^1$.*

Moreover, for $t = 3$ the algebra $R(p, \lambda)$ is a graded complete intersection of the form $k[X_1, \dots, X_s]/(\rho_3, \dots, \rho_s)$ where $s = 3$ or 4 and $\rho_3, \dots, \rho_s$ is a homogeneous regular sequence. For $k = \mathbb{C}$ the assertion also holds for $t \geq 4$ for $R(p, \lambda')$ for a suitable choice of parameters. $\lambda' = (\lambda'_3, \dots, \lambda'_t)$.

We remark that in almost every case $\text{Root } f_A(T) \subset \mathbb{S}^1$ implies that the Coxeter transformation φ_A is periodic. In fact the weight sequences $(3, 3, 3, 3)$ and $(2, 2, 2, 2, 4)$ are the only exceptions (section 4).

Theorem 4. *Let $C = C(p, \lambda)$ be a wild canonical algebra with weight sequence $p = (p_1, \dots, p_t)$. Consider $A = C[P]$ an extended canonical algebra. The following are equivalent:*

- (a) *$R(p, \lambda)$ is formally 3-generated*
- (b) *φ_A is periodic of period d and there is a primitive d -th root of unity which is root of $f_A(T)$.*

For $t = 3$ the algebra $R(p, \lambda)$ is always a graded complete intersection of the form $k[X_1, X_2, X_3]/(f)$. Moreover, for $t \geq 4$ and $k = \mathbb{C}$ this also holds for $R(p, \lambda')$ for a suitable choice of parameters $\lambda' = (\lambda'_3, \dots, \lambda'_t)$.

For the proof of Theorem 3 (resp. Theorem 4) we classify in section 5 all the weight sequences p such that $R(p, \lambda)$ is formally a complete intersection (resp. $R(p, \lambda)$ has 3 homogeneous generators). In the complex case the algebras $R(p, \lambda)$ of Theorem 3 correspond to the Fuchsian singularities which are minimal elliptic [36, Proposition 5.5.1] and the classification is related to Laufer's [20]. The algebras $R(p, \lambda)$ of Theorem 4 relate to classifications by Dolgachev [6] and Wagreich [37] and include the 14 exceptional unimodal Arnold's singularities [1]. We refer the reader to the complete account by Ebeling [9].

The research for this work was done during exchange visits México-Paderborn. We thank our universities and CONACyT in México for support. We thank Henning Krause for directing our attention to Buchweitz's paper [5].

1.2. Notation and conventions. Let Q be a finite quiver without oriented cycles. The *path algebra* kQ has as basis all the oriented paths in Q and product given by juxtaposition of paths. Given an ideal I of kQ which is *admissible* (that is, $(kQ^+)^m \subset I \subset (kQ^+)^2$ for some $m \geq 2$, where kQ^+ is the ideal of kQ generated by the arrows), we consider the finite dimensional k -algebra $A = kQ/I$. By ‘module’ we mean a finite dimensional right A -module. The category of modules is denoted $\text{mod-}A$. A module is identified with a covariant functor $X: kQ \rightarrow \text{mod-}k$ such that $X(\rho) = 0$ for every $\rho \in I$. Important modules are the simple modules S_i associated to vertices $i \in Q_0$ (= set of vertices of Q), the projective cover $P_i = A(-, i)$ of S_i and the injective envelope $I_i = DA(i, -)$ of S_i , where $D = \text{Hom}_k(-, k)$ is the canonical duality.

We denote by $K_0(A)$ the Grothendieck group of A . Since A has finite global dimension, the classes $[P_i]$ with $i \in Q_0$ form a basis of $K_0(A)$. Thus the *Coxeter transformation* $\varphi_A: K_0(A) \rightarrow K_0(A)$, given by $\varphi_A([P_i]) = -[I_i]$ defines an isomorphism. The Grothendieck group $K_0(A)$ is equipped with a bilinear form $\langle -, - \rangle_A: K_0(A) \times K_0(A) \rightarrow \mathbb{Z}$, called the *Euler form*, defined in the classes of modules X and Y as $\langle [X], [Y] \rangle_A = \sum_{i=0}^{\infty} (-1)^i \dim_k \text{Ext}_A^i(X, Y)$. In case $A = k\Delta$ is a hereditary algebra, then $\varphi_A([X]) = [\tau_A X]$ for any indecomposable non-projective A -module X . For the general situation, we have to look at the derived category $D(A) = D^b(\text{mod-}A)$ of bounded complexes of A -modules.

The *derived category* $D(A)$ contains a copy $\text{mod-}A[n]$ of $\text{mod-}A$ for each integer $n \in \mathbb{Z}$, with objects written $X[n]$ and satisfying

$$\text{Hom}_{D(A)}(X[n], Y[m]) = \text{Ext}_A^{n-m}(X, Y).$$

We say that an algebra A is *derived hereditary* (resp. *derived canonical*) if $D(A)$ is triangle equivalent to $D(H)$ (resp. $D(C)$) for a hereditary algebra H (resp. a canonical algebra C).


The category $D(A)$ has Auslander-Reiten triangles which yield a self-equivalence $\tau_{D(A)}$ of $D(A)$, the Auslander-Reiten translation, satisfying $\text{Hom}_{D(A)}(Y, \tau_{D(A)} X[1]) = D\text{Hom}_A(X, Y)$. The natural isomorphism $K_0(A) \rightarrow K_0(D(A))$, $X \mapsto X[0]$ yields $\varphi_A([X]) = [\tau_{D(A)} X]$.

For background material on representations of algebras and derived categories we refer the reader to [14, 34].

For vectors $v, w \in K_0(A)$ we get $\langle v, \varphi_A(w) \rangle_A = -\langle w, v \rangle_A$.

2. Extended canonical algebras: basic properties


2.1. Let $C = C(p, \lambda)$ be the *canonical algebra* defined by the weight sequence $p = (p_1, \dots, p_t)$ with $p_i \geq 2$ and $(\lambda = \lambda_3, \dots, \lambda_t)$ a sequence of pairwise distinct non-zero elements of k , that is, C is defined by the quiver



satisfying the $t - 2$ equations

$$\alpha_{ip_i} \dots \alpha_{i2} \alpha_{i1} = \alpha_{2p_2} \dots \alpha_{22} \alpha_{21} - \lambda_i \alpha_{1p_1} \dots \alpha_{12} \alpha_{11}, \quad i = 3, \dots, t.$$

The algebra C is a one-point extension $H[M]$ of the hereditary algebra $H = C/(0)$ by an H -module M with dimension vector $[M] = (\dim_k M(i))_i \in K_0(H)$ as follows:



In case $t \geq 3$, the module M is an indecomposable H -module which is not preprojective or preinjective.

Observe that the underlying graph of H is a star $[p_1, p_2, \dots, p_t]$ with linear arms having p_i vertices, $i = 1, \dots, t$. If H is representation-finite, then $[p_1, p_2, \dots, p_t]$ is a Dynkin diagram (that is, $\sum_{i=1}^t \frac{1}{p_i} > t - 2$) and C is *tame of domestic type*. If H is tame, then $[p_1, p_2, \dots, p_t]$ is an extended Dynkin diagram (that is, $\sum_{i=1}^t \frac{1}{p_i} = t - 2$) and C is *tame of tubular type*. See [34] for details.

2.2. The representation theory of $\text{mod-}C$ for $C = C(p, \lambda)$ a canonical algebra is controlled by the category $\text{coh}(\mathbb{X})$ of coherent sheaves on a weighted projective line $\mathbb{X} = \mathbb{X}(p, \lambda)$, since the derived categories $D(C)$ and $D^b(\text{coh}(\mathbb{X}))$ are equivalent as triangulated categories [12]. The complexity of the classification problem for $\text{coh}(\mathbb{X})$, and hence for $\text{mod-}C$, is essentially determined by the (orbifold) Euler characteristic $\chi_{\mathbb{X}} = 2 - \sum_{i=0}^t (1 - 1/p_i)$. Indeed, for $\chi_{\mathbb{X}} > 0$, the algebra C is tame of domestic type and for $\chi_{\mathbb{X}} = 0$, the algebra C is tubular. The wild case $\chi_{\mathbb{X}} < 0$ was carefully studied in [23], a paper which is the basis of the present investigation.

2.3. Let $C = C(p, \lambda)$ be a canonical algebra. Let P be an indecomposable projective or injective C -module, then $A = C[P]$ is called an *extended canonical algebra*. Hence A arises from by adjoining one arrow with a new vertex to an arbitrary vertex of C while keeping the relations for C without introducing any new relations. In particular, the opposite algebra of an extended canonical algebra is again extended canonical.

Lemma. *Any extended canonical algebra is wild.*

Proof: Recall that for an algebra $B = kQ/I$ where Q has no oriented cycles and I is generated by $\rho_1, \dots, \rho_s \in \bigcup_{i,j \in Q_0} I(i, j)$, the Tits (quadratic) form $q_B: K_0(B) \rightarrow \mathbb{Z}$ is defined by

$$q_B(x) = \sum_{i \in Q_0} x(i)^2 - \sum_{i \rightarrow j} x(i)x(j) + \sum_{i,j \in Q_0} r(i, j)x(i)x(j),$$

where $r(i, j) = \#\{s: \rho_s \in I(i, j)\}$. The Tits form is weakly non-negative (i.e. $q_A(v) \geq 0$ for $v \in \mathbb{N}^{Q_0}$) if B is a tame algebra [31].

For an extended canonical algebra $A = C[P]$ with extension vertex $*$ such that $\text{rad } P_* = P = P_j$ for some vertex j in C , we have the following:

- since $\text{gl dim } A = 2$, then $q_A(x) = \langle x, x \rangle_A$;
- the vector $w \in K_0(C) \subset K_0(A)$ with $w(i) = 1$ for every i in C , satisfies $q_c(w) = 0$;
- for $e_* = [S_*]$, we get

$$q_A(2w + e_*) = 4q_c(w) - 2w(j) + 1 < 0.$$

Hence A is of wild type. □

2.4. The following result is fundamental for introducing the concept of extended canonical algebras.

Proposition. *Let X and Y be two indecomposable projective or injective C -modules over a canonical algebra C . Then the extended canonical algebras $C[X]$ and $C[Y]$ are derived equivalent.*


In particular, the derived class of an extended canonical algebra is independent of the chosen projective module.

Proof. (See [24]). Under the equivalence $D(C) = D^b(\text{coh}(\mathbb{X}(p, \lambda)))$ the modules X and Y become line bundles over $\mathbb{X}$ (up to translation in $D^b(\text{coh}(\mathbb{X}))$). Since the Picard group of $\mathbb{X}$ acts transitively on isomorphism classes of line bundles, there is a self-equivalence of $D(C)$ sending X to Y . The assertion follows from [2]. $\square$

2.5. The following remark is useful:

Proposition. *Let A be an extended canonical algebra of $C = C(p, \lambda)$ with $p = (p_1, \dots, p_t)$ satisfying $t \geq 3$. Then A is derived equivalent to a one-point extension $H[N]$ of a hereditary algebra H by an indecomposable module N .*

Proof. By (2.3), we may assume that A is the path algebra of the quiver



equipped with the canonical relations $\alpha_{ip_i} \dots \alpha_{i1} = \alpha_{2p_2} \dots \alpha_{21} - \lambda_i \alpha_{1p_1} \dots \alpha_{11}$ ($3 \leq i \leq t$). It follows that A is the one-point extension of the path algebra of the star $[2, p_1, p_2, \dots, p_t]$ by an indecomposable module $N = N(\lambda_3, \dots, \lambda_t)$ whose restriction to $[p_1, p_2, \dots, p_t]$ is M as in (2.1) and $N(*) = 0$. $\square$

Remark: An obvious variant of the above statement yields families of one-point extensions of hereditary algebras which are pairwise derived equivalent. For instance, for the canonical type $(2, 3, 7)$, the Proposition yields $10 = (2-1) + (3-1) + (7-1) + 1$ choices of pairs (H, N) of a hereditary algebra H and an indecomposable H -module N such that $H[N]$ is derived equivalent to an extended canonical algebra of type $(2, 3, 7)$.

3. The derived category of an extended canonical algebra

3.1. In this section we are going to investigate the nature of the bounded derived category of an extended canonical algebra $A = C[P]$. As it turns out the structure of this triangulated category will sensibly depend on the sign of the (orbifold) Euler characteristic $\chi_{\mathbb{X}}$ of the weighted projective line $\mathbb{X}$ associated to C .

Let $\mathcal{T}$ be a triangulated k -category, see [13, 27, 18] for definition and properties. An object E in $\mathcal{T}$ is called *exceptional* if $\text{End}(E) = k$ and $\text{Hom}(E, E[n]) = 0$ for all integers $n \neq 0$. By ${}^{\perp}E$ (resp. $E^{\perp}$) we denote the full triangulated subcategory of $\mathcal{T}$ consisting of all objects $X \in \mathcal{T}$ (resp. $Y \in \mathcal{T}$) satisfying $\text{Hom}(X, E[n]) = 0$ (resp. $\text{Hom}(E[n], Y) = 0$ for each integer n). By [3] the inclusion ${}^{\perp}E \hookrightarrow \mathcal{T}$ (resp. $E^{\perp} \hookrightarrow \mathcal{T}$) admits an exact left (resp. right) adjoint $\ell : \mathcal{T} \rightarrow {}^{\perp}E$ (resp. $r : \mathcal{T} \rightarrow E^{\perp}$).

For the purpose of this paper we call an exceptional object *special in $\mathcal{T}$* if one of the following two conditions is satisfied:

- (i) the left perpendicular category ${}^{\perp}E$ is equivalent to $D^b(\text{coh}(\mathbb{X}))$ for some weighted projective line $\mathbb{X}$ and, moreover, the left adjoint ℓ maps E to a line bundle in $\text{coh}(\mathbb{X})$.
- (ii) the right perpendicular category $E^{\perp}$ is equivalent to $D^b(\text{coh}(\mathbb{X}))$ for some weighted projective line $\mathbb{X}$ and, moreover, the right adjoint r maps E to a line bundle in $\text{coh}(\mathbb{X})$.

Again, for the purpose of this paper, an object T of $\mathcal{T}$ is called a *tilting object* in $\mathcal{T}$ if (i) T generates $\mathcal{T}$ as a triangulated category, (ii) $\text{Hom}(T, T[n]) = 0$ holds for each non-zero integer n , and (iii) the endomorphism algebra of T has finite global dimension.

3.2. Our main tool to investigate the shape of $D(A)$ is the following proposition.

Proposition. *Let $\mathcal{T}$ be a triangulated category having an exceptional object E that is special in $\mathcal{T}$. Then there exists a tilting object $\bar{T}$ of $\mathcal{T}$ whose endomorphism ring is an extended canonical algebra.*

Proof. With the previous notations we assume that $E^\perp = D^b(\text{coh}(\mathbb{X}))$ and $r(E)$ is a line bundle in $\text{coh}(\mathbb{X})$. (The assumption on the left perpendicular ${}^\perp E$ category is treated similarly.) We choose a tilting object T in $\text{coh}(\mathbb{X})$, hence in $D^b(\text{coh}(\mathbb{X}))$ having the line bundle $r(E)$ as a direct summand and such that $\text{End}(T) = C$ is the canonical algebra attached to $\mathbb{X}$ (see [11]). We claim that $\bar{T} = T \oplus E$ is a tilting object in $\mathcal{T}$. Indeed, since T generates $D^b(\text{coh} \mathbb{X})$ and E together with $E^\perp$ generates $\mathcal{T}$, it follows that $\bar{T}$ generates $\mathcal{T}$. Next, we show that $\text{Hom}(\bar{T}, \bar{T}[n]) = 0$ for each nonzero integer n . This reduces to show that $\text{Hom}(E[n], T) = 0$ and $\text{Hom}(T, E[n])$ holds for every nonzero n . The first assertion holds for each n since T belongs to $E^\perp$. For the second we use that $\text{Hom}(T, E[n]) = \text{Hom}(T[-n], r(E))$ is zero since by construction $r(E)$ is a direct summand of the tilting object T . Finally, the endomorphism ring of $\bar{T}$ is given as the matrix ring

$$\begin{pmatrix} C & 0 \\ P & k \end{pmatrix}, \quad \text{where } P = \text{Hom}(T, E) = \text{Hom}(T, rE)$$

is an indecomposable projective C -module, hence $\text{End}(\bar{T}) = C[P]$ is an extended canonical algebra. Moreover, as it is easily seen, $C[P]$ has global dimension two. $\square$

Keeping the assumptions on E and $\mathcal{T}$ from the proposition, we obtain.





Corollary. *If $\mathcal{T}$ is triangle equivalent to a bounded derived category $D(B)$ for some finite dimensional k -algebra B or, more generally, if $\mathcal{T}$ is algebraic in the sense of Keller [18], then $\mathcal{T}$ is triangle equivalent to $D^b(\text{mod-}A)$, for the extended canonical algebra $A = C[P]$.*

Proof. The first claim follows from [32], the general version requires [18] or [4]. $\square$

3.3. Positive Euler characteristic: the domestic case. Consider a canonical algebra $C = C(p_1, \dots, p_t)$ of domestic type, that is, $\sum_{i=1}^t \frac{1}{p_i} > 1$. Let $\Delta = [p_1, \dots, p_t]$ be the star corresponding to the weight sequence $p = (p_1, \dots, p_t)$ and $\tilde{\Delta}$ be the corresponding extended Dynkin diagram. Then $\tilde{\Delta}$ admits a unique positive *additive function* λ assuming value 1, that is, $\lambda: \tilde{\Delta} \rightarrow \mathbb{N}$ satisfies the conditions:

- (i) $2\lambda(i) = \sum_{j \in i^+} \lambda(j)$, where i^+ is the set of neighbors of i ;
- (ii) for some vertex $i \in \tilde{\Delta}_0$, $\lambda(i) = 1$. Such an i is called an *extension vertex*.

The *double extended graph* of type Δ , denoted by $\tilde{\tilde{\Delta}}$, is the graph arising from $\tilde{\Delta}$ by adjoining a new edge in an extension vertex. The list of double extended Dynkin graphs is the following:

weight sequence	double extended Dynkin graph
(p, q)	
$(2, 2, 4)$	
$(2, 3, 3)$	 $(= [3, 3, 4])$
$(2, 3, 4)$	 $(= [2, 4, 5])$
$(2, 3, 5)$	 $(= [2, 3, 7])$

Proposition. *Let $C = C(p, \lambda)$ be a canonical algebra of domestic type $p = (p_1, p_2, p_3)$. Let $\Delta = [p_1, p_2, p_3]$ be the associated Dynkin diagram. For any indecomposable pre-projective C -module N , the one-point extension $A = C[N]$ is derived equivalent to a hereditary algebra of type $\tilde{\tilde{\Delta}}$. In particular, an extended canonical algebra of weight type p is derived hereditary of type $\tilde{\tilde{\Delta}}$.*

Proof. The algebra C is tilted of a hereditary algebra $H = k\tilde{\Delta}_1$, where $\tilde{\Delta}_1$ is a quiver with underlying graph $\tilde{\Delta}$. There is a derived equivalence $F: D^b(\text{mod-}H) \rightarrow D^b(\text{mod-}C)$ sending the indecomposable projective P_i corresponding to an extension vertex i of $\tilde{\Delta}$ into the projective C -module P_ω . Observe that $H[P_i] = k\tilde{\Delta}_1$ is a hereditary algebra where $\tilde{\Delta}_1$ is a quiver with underlying graph $\tilde{\Delta}$.

Let $\mathbb{X} = \mathbb{X}(p, \lambda)$ be a weighted projective line such that $D^b(\text{mod-}C) = D^b(\text{coh}(\mathbb{X}))$. As an object in $\text{coh}(\mathbb{X})$ the object P_ω has rank one (see [11]). Also by [11], there is an equivalence in $D^b(\text{coh}(\mathbb{X}))$ sending P_ω to any indecomposable preprojective C -module N . By [2], the one-point extension $C[N]$ is derived equivalent to $C[P_\omega]$ which is derived hereditary of type $\tilde{\Delta}$. $\square$

3.4. The converse of Proposition 3.3 also holds.

Proposition. *Let $C = C(p, \lambda)$ be a canonical algebra and P be an indecomposable projective C -module. The extended canonical algebra $A = C[P]$ is derived hereditary if and only if C is tame domestic.*

Proof. If C is tame domestic, then A is derived hereditary by (2.5). For the converse, consider the set of weight sequences $p = (p_1, p_2, \dots, p_t)$ with $2 \leq p_1 \leq p_2 \leq \dots \leq p_t$ with the domination order defined in (1.1).

The statement follows by induction on the domination order from the following two facts:

- (a) a canonical tubular algebra C is not derived hereditary;
- (b) any wild weight sequence dominates a tubular one;
- (c) if M is an indecomposable B -module such that $B[M]$ is derived hereditary, then B is derived hereditary.

(a): follows from the structure of derived categories of hereditary algebras, see [14].
 (b) is clear.

(c): Assume $D^b(\text{mod-}B[M]) = D^b(\text{mod-}H)$ for a hereditary algebra H . By [12], $D^b(\text{mod-}B)$ is equivalent to the right perpendicular category in $D^b(\text{mod-}H)$ with respect to an exceptional object E , that is, E is an indecomposable object satisfying $\text{Ext}^1(E, E) = 0$ and

$$D^b(\text{mod-}B) = E^\perp = \{X \in D^b(\text{mod-}H) : \text{Hom}(E, X) = 0 = \text{Ext}^1(E, X)\}.$$

Without loss of generality we may assume that $E \in \text{mod-}H$. Then $D^b(\text{mod-}B) \cong D^b(E^\perp)$, where now $E^\perp$ is formed in $\text{mod-}H$. Hence $E^\perp = \text{mod-}H'$ for a hereditary algebra H' . $\square$

3.5. Euler characteristic zero: the tubular case. Consider a canonical algebra $C = C(p, \lambda)$ with weight sequence $p = (p_1, \dots, p_t)$, we shall assume that $2 \leq p_1 \leq p_2 \leq \dots \leq p_t$. The module category $\text{mod-}C$ accepts a *separating tubular family* $\mathcal{T} = (T_\lambda)_{\lambda \in \mathbb{P}_1 k}$, where T_λ is a homogeneous tube for all λ with the exception of t tubes $T_{\lambda_1}, \dots, T_{\lambda_t}$ with T_{λ_i} of rank p_i ($1 \leq i \leq t$). See [34].

Let $\mathbb{X} = \mathbb{X}(p, \lambda)$ be the weighted projective line such that $\text{mod-}C$ and $\text{coh}(\mathbb{X})$ are derived equivalent. We fix an equivalence $D^b(\text{mod-}C) = D^b(\text{coh}(\mathbb{X}))$. Let S be a simple C -module in the mouth of the tube of rank p_t and consider S as an object in $\text{coh}(\mathbb{X})$. The category $S^\perp$ right perpendicular to the object S is the full subcategory of $\text{coh}(\mathbb{X})$ consisting of all $\mathcal{F} \in \text{coh}(\mathbb{X})$ satisfying

$$\text{Hom}_{\mathbb{X}}(S, \mathcal{F}) = 0 = \text{Ext}_{\mathbb{X}}^1(S, \mathcal{F}).$$

By [12], $S^\perp = \text{coh}(\mathbb{X}')$ where $\mathbb{X}' = \mathbb{X}(p', \lambda)$ is a weighted projective line with weight sequence $p' = (p_1, p_2, \dots, p_{t-1}, p_t - 1)$. Moreover, if $0 \rightarrow \tau S \rightarrow U \rightarrow S \rightarrow 0$ is the almost split sequence in $\text{coh}(\mathbb{X})$, then U is a simple object in $S^\perp$ of τ' -period $p_t - 1$, where $\tau' = \tau_{D^b(\text{coh}(\mathbb{X}'))}$.

Proposition. *Let $C = C(p, \lambda)$ be a canonical algebra of tubular type $p = (p_1, \dots, p_t)$ and $A = C[P]$ be an extended canonical algebra. Then A is derived canonical of type $\bar{p} = (p_1, \dots, p_{t-1}, p_t + 1)$.*

Proof. By (2.4), we may choose P to be the simple projective C -module. We shall show that A is quasi-tilted of type $\bar{p} = (p_1, \dots, p_{t-1}, p_t + 1)$, see [15].

Let $\mathbb{X}$ be a weighted projective line with $D^b(\text{coh}(\mathbb{X})) = D^b(\text{mod-}C)$ and let $\bar{\mathbb{X}}$ denote a weighted projective line of type $\bar{p}$ such that $\text{coh}(\mathbb{X})$ is the perpendicular category $S^\perp$ formed in $\text{coh}(\bar{\mathbb{X}})$ for a simple S from the tube of rank $p_t + 1$. Let U be the middle term of the almost split sequence $0 \rightarrow \tau S \rightarrow U \rightarrow S \rightarrow 0$ in $\text{coh}(\bar{\mathbb{X}})$. Then U is a simple in $S^\perp = \text{coh}(\mathbb{X})$ belonging to the largest tube in $\text{coh}(\mathbb{X})$.

By hypothesis, $\mathbb{X}$ has tubular type. By [22], there is a tilting object $\mathcal{T}$ in $\text{coh}_0(\mathbb{X}) \vee \text{coh}_+(\mathbb{X})[1]$ such that $\text{End}(\mathcal{T}) = C$, where $\text{coh}_0(\mathbb{X})$ (resp. $\text{coh}_+(\mathbb{X})$) denotes the full

subcategory of $\text{coh}(\mathbb{X})$ formed by the sheaves of rank 0 (resp. positive rank). Therefore, $\mathcal{T} \cup \{S\} \subset \text{coh}_0(\bar{\mathbb{X}}) \vee \text{coh}_+(\bar{\mathbb{X}})[1]$ is a tilting complex for $\bar{\mathbb{X}}$ whose endomorphism ring is isomorphic to $C[P] = A$. $\square$

3.6. Negative Euler characteristic: the wild case. For negative Euler characteristic the derived category of modules over an extended canonical algebra $C[P]$ relates to the study of the $\mathbb{Z}$ -graded surface singularity R associated with C and the weighted projective line $\mathbb{X}$ associated to C . We refer to [11, 21, 12] for further details. The weighted projective line $\mathbb{X} = \mathbb{X}(p, \lambda)$ for a weight sequence $p = (p_1, \dots, p_t)$ and a parameter sequence $\lambda = (\lambda_3, \dots, \lambda_t)$ was introduced in [11] by means of the algebra

$$S = S(p, \lambda) := k[x_1, \dots, x_t] / (x_i^{p_i} = x_2^{p_2} - \lambda_i x_1^{p_1}), \quad i = 3, \dots, t.$$

The algebra S is naturally graded over the abelian group $\mathbb{L}(p)$ with generators $\vec{x}_1, \dots, \vec{x}_t$ and relations $p_1 \vec{x}_1 = \dots = p_t \vec{x}_t =: \vec{c}$ by giving each x_i the degree $\vec{x}_i$. Here, $\vec{c}$ is called the canonical element of $\mathbb{L}(p)$. The group $\mathbb{L}(p)$ is isomorphic to the direct sum of the group $\mathbb{Z}$ of integers and some finite group. A quick way to arrive at the category $\text{coh}(\mathbb{X})$ of coherent sheaves on $\mathbb{X}$ is by putting

$$\text{coh}(\mathbb{X}) = \frac{\text{mod}^{\mathbb{L}(p)}\text{-}S}{\text{mod}_0^{\mathbb{L}(p)}\text{-}S},$$

where the quotient category on the right is formed in the sense of [10] and the categories $\text{mod}^{\mathbb{L}(p)}\text{-}S$ (resp. $\text{mod}_0^{\mathbb{L}(p)}\text{-}S$) are the categories of finitely generated $\mathbb{L}(p)$ -graded S -modules (resp. those of finite length).

The element $\vec{\omega} = (t-2)\vec{c} - \sum_{i=0}^t \vec{x}_i$ from $\mathbb{L}(p)$ is called the *dualizing element*. Its importance comes from the fact that Serre duality for $\text{coh}(\mathbb{X})$ holds in the form $D\text{Ext}^1(X, Y) = \text{Hom}(Y, X(\vec{\omega}))$, where $X \mapsto X(\vec{\omega})$ is the self-equivalence of $\text{coh}(\mathbb{X})$ induced by grading shift $M \mapsto M(\vec{\omega})$, given by $M(\vec{\omega})_{\vec{x}} = M_{\vec{\omega}+\vec{x}}$.

Assume that $\chi_{\mathbb{X}} < 0$. Then the *graded surface singularity* $R = R(p, \lambda)$ attached to the weighted projective line $\mathbb{X}$ (or the canonical algebra C) with data (p, λ) is defined as $R = \bigoplus_{n=0}^{\infty} R_n$, where $R_n = S_{n\vec{\omega}}$. It follows immediately that R is a finitely generated, i.e. affine, k -algebra where each R_n is finite dimensional over k and, moreover, $R_0 = k$ and $R_1 = 0$. The next theorem illustrates the role of R , and shows in particular that the algebra R keeps all information on $\mathbb{X}$. For the proofs we refer to [21, 12].

Theorem. *Assume $\chi_{\mathbb{X}} < 0$. Then the following holds:*

- (a) *The algebra $R = R(p, \lambda)$ is a positively $\mathbb{Z}$ -graded isolated surface singularity which is graded Gorenstein of Gorenstein index -1 .*
- (b) *There is a natural equivalence $\mathrm{coh}(\mathbb{X}) \rightarrow \mathrm{mod}^{\mathbb{Z}}\text{-}R/\mathrm{mod}_0^{\mathbb{Z}}\text{-}R$, induced by restricting the grading from $\mathbb{L}(p)$ to $\mathbb{Z} = \mathbb{Z}\vec{\omega}$.*
- (c) *For $k = \mathbb{C}$, the algebra $R(p, \lambda)$ is the positively $\mathbb{Z}$ -graded algebra of automorphic forms on the (upper) complex half-plane $\mathbb{H}_+$ with respect to the action of a Fuchsian group G of the first kind of signature $(0; p_1, \dots, p_t)$.* $\square$

Concerning (a) we note that — restricting to the case of Krull dimension two — the Gorenstein index d can be defined through the minimal graded injective resolution $0 \rightarrow R \rightarrow E^0 \rightarrow E^1 \rightarrow E^2 \rightarrow 0$ of the R -module R , where the term E^2 is the graded injective hull of $k(d)$ and generally the *grading shift* (n) is defined by $M(n)_m = M_{n+m}$. In this situation, Serre duality holds for $\mathrm{coh}(\mathbb{X})$ in the form $D\mathrm{Ext}^1(X, Y) = \mathrm{Hom}(Y, X(-d))$, such that the Auslander-Reiten translation comes from the grading shift $X \mapsto X(-d)$.

Concerning (c) we remark that G is the orbifold fundamental group of $\mathbb{X}$, having a presentation $\langle \sigma_1, \dots, \sigma_t \mid \sigma_1^{p_1} = \dots = \sigma_t^{p_t} = \sigma_1 \cdots \sigma_t \rangle$ acting by covering transformations on the (branched) universal cover $\mathbb{H}_+$ of $\mathbb{X}$. We refer to [26, 28] for the associated rings of automorphic forms.

3.7. For a variety X Orlov investigated in [29] the triangulated category $D_{\mathrm{Sg}}(X)$ of the singularities of X defined as the quotient of the bounded derived category $D^b(\mathrm{coh}(X))$ of coherent sheaves modulo the full subcategory of perfect complexes. If X is affine with coordinate algebra R this category $D_{\mathrm{Sg}}(R)$ is just the quotient $D^b(\mathrm{mod}\text{-}R)/D^b(\mathrm{proj}\text{-}R)$, where $\mathrm{proj}\text{-}R$ is the category of finitely generated projective R -modules. In [30] Orlov further introduced a graded variant

$$D_{\mathrm{Sg}}^{\mathbb{Z}}(R) = D^b(\mathrm{mod}^{\mathbb{Z}}\text{-}R)/D^b(\mathrm{proj}^{\mathbb{Z}}\text{-}R)$$

called the *triangulated category of the graded singularity* R which will play a central role in this section.

Under the name *stabilized derived category of R* the categories $D_{\mathrm{Sg}}(R)$ were introduced by Buchweitz in [5]. His results easily extend to the graded case and yields for an R that is graded Gorenstein an alternative description of $D_{\mathrm{Sg}}^{\mathbb{Z}}(R)$ as the *stable*

category of graded maximal Cohen-Macaulay modules $\underline{\text{MCM}}^{\mathbb{Z}}\text{-}R$. More precisely, he showed that the category $\text{MCM}^{\mathbb{Z}}\text{-}R$ of maximal graded Cohen-Macaulay R -modules is a Frobenius-category, hence inducing — in Keller's terminology [18] — on the attached stable category $\underline{\text{MCM}}^{\mathbb{Z}}\text{-}R$ of graded maximal Cohen-Macaulay modules modulo projectives, the structure of an *algebraic* triangulated category. For a related approach measuring the complexity of a singularity by a triangulated category we refer to Krause's account [19].

Let $R = \bigoplus_{n \geq 0} R_n$, $R_n = S_{n\bar{\omega}}$, be the positively $\mathbb{Z}$ -graded Gorenstein singularity attached to the weighted projective line $\mathbb{X}$. It follows from [21, 5.6] that R has Krull dimension two and Gorenstein index -1 . We fix some notation: Let $\mathcal{M} = \text{D}^b(\text{mod}^{\mathbb{Z}}\text{-}R)$ and $\mathcal{M}_+ = \text{D}^b(\text{mod}^{\mathbb{Z}_+}\text{-}R)$. Let $\mathcal{P}_+$ be the triangulated subcategory of $\mathcal{M}_+$ generated by all $R(-n)$, $n \geq 0$ and $\mathcal{T}$ its left perpendicular category ${}^{\perp}\mathcal{P}_+$ formed in $\mathcal{M}_+$. Denote further by $\mathcal{S}_+$ the triangulated subcategory of $\mathcal{M}_+$ generated by all $k(-n)$, $n \geq 0$ and $\mathcal{D}$ its right perpendicular category $\mathcal{S}_+^{\perp}$ formed in $\mathcal{M}_+$. Finally let $\mathcal{D}(-1) = \mathcal{S}_+(-1)^{\perp}$. Then [30, 2.5] implies the following proposition.

Proposition. *Assume that $\chi_{\mathbb{X}} < 0$ and let R be the positively $\mathbb{Z}$ -graded singularity attached to $\mathbb{X}$. Then the following holds:*

(a) *The natural functor $\mathcal{T} \hookrightarrow \mathcal{M} \xrightarrow{q} \text{D}_{\text{Sg}}(R)$, where q is the quotient functor, is an equivalence of triangulated categories.*

(b) *The R -module k is an exceptional object in $\mathcal{T}$ with ${}^{\perp}k = \mathcal{D}(-1)$. Moreover, the category $\text{D}^b(\text{coh}(\mathbb{X}))$ is naturally equivalent to $\mathcal{D}(-1)$ under the functor $Y \mapsto (\mathbf{R}\Gamma_+(Y))(-1)$.*

Proof. For the convenience of the reader we sketch the argument. Using that R is Gorenstein of Gorenstein index -1 , and invoking Gorenstein duality $\mathbf{R}\text{Hom}_R^{\bullet}(-, R)$ of $\mathcal{M}$ one sees that $T^{\perp} \subset \mathcal{D}(-1)^{\perp}$ and hence $\mathcal{D}(-1)$ is a full subcategory of $\mathcal{T}$. Further we see that ${}^{\perp}k = \mathcal{D}(-1)$. It is well-known that

$$\Gamma_+ : \text{coh}(\mathbb{X}) \rightarrow \text{mod}^{\mathbb{Z}_+}\text{-}R, \quad Y \mapsto \bigoplus_{n=0}^{\infty} \text{Hom}(\mathcal{O}, Y(n))$$

is a full embedding having sheafification, that is, the quotient functor $q_+ : \text{mod}^{\mathbb{Z}_+}\text{-}R \rightarrow \text{coh}(\mathbb{X})$ as an exact left adjoint and such that composition $q\Gamma_+$ is the identity functor on $\text{coh}(\mathbb{X})$, compare [11, 1.8], [21, 5.7]. It follows that $\mathbf{R}\Gamma_+ : \text{D}^b(\text{coh}(\mathbb{X})) \rightarrow \mathcal{M}_+$ is a full embedding having $q_+ : \mathcal{M}_+ \rightarrow \mathcal{M}_+/\mathcal{S}_+$ as a left adjoint, and $q_+\mathbf{R}\Gamma_+ = 1$.

Since R is positively graded with $R_0 = k$, it follows that k is exceptional in $\mathcal{M}$ and hence in $\mathcal{M}_+$. Invoking the minimal graded injective resolution $0 \rightarrow R \rightarrow E^0 \rightarrow E^1 \rightarrow E^2 \rightarrow 0$, where E^0 and E^1 are socle-free and E^2 is the graded injective hull of $k(-1)$, it follows that k belongs to $\mathcal{T}$ and then also to $\mathcal{D}(-1)$. It is straightforward to check that $\mathcal{D}(-1)^\perp$ equals the triangulated subcategory $\langle k \rangle$ generated by k , and hence ${}^\perp k = \mathcal{D}(-1)$ in $\mathcal{T}$. $\square$

3.8. Proof of Theorem 1. We are now in a position to clarify the structure of the category $D_{\text{Sg}}^{\mathbb{Z}}(R)$. The result was first observed by K. Saito and A. Takahashi (personal communication); it is not yet published, and uses the technique of matrix factorizations as in [17].

By Proposition 3.2 it suffices to show that the left adjoint $\ell : \mathcal{T} \rightarrow {}^\perp k$ to the inclusion $j : {}^\perp k \hookrightarrow \mathcal{T}$ maps k to a line bundle in $\mathcal{D}(-1) = D^b(\text{coh}(\mathbb{X}))$ up to translation in $\mathcal{D}(-1)$. We put $A = (\mathbf{R}\Gamma_+(\mathcal{O}(-\vec{\omega})))(-1)$ and construct a morphism $\gamma : k \rightarrow A[1]$ such that $\text{Hom}(\gamma, Y) : \text{Hom}(A[1], Y) \rightarrow \text{Hom}(k, Y)$ is an isomorphism for each $Y \in {}^\perp k$ such that $\ell(k) = A[1]$.

The claim is proved in two steps. Put $R_+ = \bigoplus_{n \geq 1} R_n$, then the exact sequence $0 \rightarrow R_+ \rightarrow R \rightarrow k$ yields an exact triangle $R \rightarrow k \xrightarrow{\alpha} R_+[1]$ in $\mathcal{M}$, where $\text{Hom}(\alpha, Y)$ is an isomorphism for each $Y \in {}^\perp k$. Note for this that R belongs to ${}^\perp \mathcal{D}(-1)$.

For the next step it is useful to identify the derived category $\mathcal{M}_+$ with the full subcategory of $D^b(\text{Mod}^{\mathbb{Z}+} R)$ consisting of all complexes with cohomology in $\text{mod}^{\mathbb{Z}+} R$. Here, $\text{Mod}^{\mathbb{Z}+} R$ denotes the category of *all* graded R -modules. Let $0 \rightarrow R(-1) \rightarrow E^0 \rightarrow E^1 \rightarrow E^2 \rightarrow 0$ be the minimal graded injective resolution of $R(1)$ such that E^2 equals the graded injective envelope of k . (This uses that R has Gorenstein index -1 .) Sheafification yields the minimal injective resolution $0 \rightarrow \mathcal{O}(\vec{\omega}) \rightarrow \tilde{E}^0 \rightarrow \tilde{E}^1 \rightarrow 0$ of $\mathcal{O}(\vec{\omega})$. Accordingly $\mathbf{R}\Gamma_+(\mathcal{O}(\vec{\omega}))$ is given by the complex

$$A : \quad \cdots \rightarrow 0 \rightarrow E_+^0(-1) \rightarrow E_+^2(-1) \rightarrow 0 \cdots, \text{ where } E_+^i = \bigoplus_{n \geq 0} E_n^i,$$

whose cohomology is concentrated in degrees zero and one and given by

$$H^0(A) = R_+, \quad H^1(A) = k(-1).$$

It follows the existence of an exact triangle

$$k(-1)[-2] \rightarrow R_+ \xrightarrow{\beta} A \rightarrow k(-1)[-1],$$


in $\mathcal{M}_+$ where, by construction, A belongs to $\mathcal{D}(-1)$. For Y from $\mathcal{D}(-1)$ we have, in particular, that Y belongs to ${}^\perp k(-1)$ implying that $\text{Hom}(\beta, Y)$ is an isomorphism. To summarize: The morphism $\gamma = [k \xrightarrow{\alpha} R_+[1] \xrightarrow{\beta[1]} A[1]]$ yields isomorphisms $\text{Hom}(\gamma, Y)$ for each $Y \in \mathcal{D}(-1)$. Hence $\ell(k)[-1] = A = \mathbf{R}\Gamma_+(\mathcal{O}(\vec{\omega}))(-1)$ is a line bundle, as claimed. $\square$

3.9. The Coxeter-Dynkin algebras of a singularity. In the theory of singularities the attached Coxeter-Dynkin diagrams, see for instance [8, 7], play an important role, in particular, since they establish a link to Lie theory.

Definition. Let k be an algebraically closed field, and $R = R(p, \lambda)$ be the $\mathbb{Z}$ -graded singularity attached to the weighted projective line $\mathbb{X}(p, \lambda)$.

(a) By the *Coxeter-Dynkin algebra of hereditary type* we mean the path algebra $D[p]$ of the hereditary star $[p_1, \dots, p_t]$ having a unique sink.

(b) By the *Coxeter-Dynkin algebra of canonical type* we mean the algebra $D(p, \lambda)$ given in terms of the quiver



with the two relations $\sum_{i=2}^t \alpha_i \beta_i = 0$ and $\alpha_1 \beta_1 = \sum_{i=3}^t \lambda_i \alpha_i \beta_i$.

(c) By the *Coxeter-Dynkin algebra of extended-canonical type* we mean the one-point extension $\hat{D}(p, \lambda)$ of the Coxeter-Dynkin algebra $D(p, \lambda)$ of canonical type above, introducing a new arrow at the sink vertex and keeping the relations.

The link to singularity theory is given by the following well known result, compare [7, 8].

Theorem. For $k = \mathbb{C}$ the Coxeter-Dynkin diagram of the singularity $R(p, \lambda)$ is the underlying digraph of the Coxeter-Dynkin algebra

(a) of hereditary type, if $\chi_{\mathbb{X}} > 0$, and then $[p_1, \dots, p_t]$ is Dynkin.

(b) of canonical type, if $\chi_{\mathbb{X}} = 0$, and then p is tubular.

(c) of extended canonical type, if $\chi_{\mathbb{X}} < 0$. □

Recall that the digraph of a finite dimensional algebra A has the underlying graph of the quiver of A as solid edges and the minimal number of relations between vertices i and j as dotted edges.

Proposition. *Assume the number of weights > 1 is at least two. Then the following holds:*

(a) *The Coxeter-Dynkin algebra $D(p, \lambda)$ of canonical type is derived equivalent to the canonical algebra $C(p, \lambda)$.*

(b) *The Coxeter-Dynkin algebra $\hat{D}(p, \lambda)$ of extended canonical type is derived equivalent to the extended canonical algebra $A = C[P]$, where $C = C(p, \lambda)$.*

Proof. Let $\mathbb{X}$ be the weighted projective line with t weighted points $x_1, \dots, x_t$ of weight $p_1, \dots, p_t$, respectively. For each $i = 1, \dots, t$ denote by U_i the unique indecomposable sheaf of length $p_i - 1$ concentrated in x_i such there exists an epimorphism $\mathcal{O} \rightarrow U_i$ in $\text{coh}(\mathbb{X})$, where $\mathcal{O}$ is the structure sheaf on $\mathbb{X}$. Moreover, let

$$S_i = U_i^{(1)} \subset U_i^{(2)} \subset \dots \subset U_i^{(p_i-1)} = U_i$$

be the complete system of subobjects of U_i . It is straightforward to verify that the object

$$T = \mathcal{O}(\vec{c}) \oplus \bigoplus_{i=1}^t \left(U_i^{(1)} \oplus U_i^{(2)} \oplus \dots \oplus U_i^{(p_i-1)} \right) \oplus \mathcal{O}(-\vec{\omega})[1]$$

is a tilting object in $D^b(\text{coh}(\mathbb{X}))$. Moreover, it is not difficult to see that the endomorphism ring of T is isomorphic to the Coxeter-Dynkin algebra $D(p, \lambda)$.

This proves (a). Assertion (b) follows from (a) noting that $\mathcal{O}(-\vec{\omega})$ is a line bundle by applying [2] or arguing as in Proposition 3.2. □

3.10. The triangulated category of singularities for nonnegative Euler characteristic. In this paper we mainly concentrate on the case $\chi_{\mathbb{X}} < 0$. To complete the picture we review the situation for $\chi_{\mathbb{X}} \geq 0$.

Assume that $k = \mathbb{C}$. For $\chi_{\mathbb{X}} > 0$ the Dynkin diagram given by the weight type and the Coxeter-Dynkin diagram of the singularity $R = R(p, \lambda)$ agree. Moreover, it is shown in [17] that the category $D_{\text{Sg}}^{\mathbb{Z}}(R)$ is equivalent to the bounded derived category $D^b(k\Delta)$ of the path algebra of a quiver of the same Dynkin type.

Next we deal with the case $\chi_{\mathbb{X}} = 0$. First, there is no $\mathbb{Z}$ -graded Gorenstein algebra R such that $\mathcal{C} = \text{mod}^{\mathbb{Z}}\text{-}R/\text{mod}_0^{\mathbb{Z}}\text{-}R$ is equivalent to $\text{coh}(\mathbb{X})$. Assume, indeed, that such an algebra R would exist. Let d denote its Gorenstein index. Since the Auslander-Reiten translation $\tau(X) = X(-d)$ has finite order 2, 3, 4 or 6 for any weighted projective line $\mathbb{X}$ of tubular type, it follows that $d = 0$, hence τ is the identity, contradiction.

Hence for $\chi_{\mathbb{X}} = 0$ it is more natural to investigate the $\mathbb{L}(p)$ -graded singularity $S = S(p, \lambda)$ and its triangulated category of singularities $\text{D}_{\text{Sg}}^{\mathbb{L}(p)}(S)$ which in the tubular case is equivalent to $\text{D}^b(\text{coh}(\mathbb{X}))$ due to recent work of Ueda [35].

4. The Coxeter polynomial of an extended canonical algebra

4.1. Let A be a finite dimensional k -algebra of finite global dimension. The *Coxeter transformation* $\varphi_A: K_0(A) \rightarrow K_0(A)$ is the automorphism induced by the Auslander-Reiten translation $\tau_{\text{D}^b(\text{mod } A)}: \text{D}^b(\text{mod } A) \rightarrow \text{D}^b(\text{mod } A)$. We shall consider φ_A as a $n \times n$ integral matrix where n is the rank of $K_0(A)$. From the Introduction, we recall that $f_A(T) = \det(T\text{id} - \varphi_A)$ is called the *Coxeter polynomial* of A .

For a one-point extension $A = B[P]$ of an algebra B by an indecomposable projective a simple calculation shows (see [24, 34]):

$$f_A(T) = (1 + T)f_B(T) - Tf_C(T).$$

Of particular interest is the hereditary case where any algebra can be constructed by repeated one-point extensions using indecomposable projective modules. In fact, for a star H of type $[p_1, \dots, p_t]$ the above formula yields:

$$f_{[p_1, \dots, p_t]}(T) := f_H(T) = \left[(T + 1) - T \sum_{i=1}^t \frac{v_{p_i} - 1}{v_{p_i}} \right] \prod_{i=1}^t v_{p_i}$$

where we set $v_n(T) = \frac{T^n - 1}{T - 1} = \sum_{i=0}^{n-1} T^i$.

For the canonical algebra $C = C(p, \lambda)$ of type $p = (p_1, \dots, p_t)$, we get

$$f_{(p_1, \dots, p_t)}(T) := f_C(T) = (T - 1)^2 \prod_{i=1}^t v_{p_i}(T).$$

In particular, all the eigenvalues of φ_C lie on the unit circle $\mathbb{S}^1$. For these calculations we refer the reader to [23].

Lemma [24]. *An extended canonical algebra $A = C[P]$ where $C = C(p, \lambda)$ with $p = (p_1, \dots, p_t)$ has Coxeter polynomial:*

$$\hat{f}_{(p_1, \dots, p_t)}(T) := f_A(T) = (T+1)(T-1)^2 \prod_{i=1}^t v_{p_i}(T) - T f_{[p_1, \dots, p_t]}(T) \quad \square$$

4.2. For later use we recall some facts on *cyclotomic polynomials*.

The n -cyclotomic polynomial $\phi_n(T)$ is inductively defined by the formula

$$T^n - 1 = \prod_{d|n} \phi_d(T).$$

Recall that the *Möbius function* is defined as follows:

$$\mu(n) = \begin{cases} 0 & \text{if } n \text{ is divisible by a square} \\ (-1)^r & \text{if } n = p_1 \dots p_r \text{ is a factorization into distinct primes.} \end{cases}$$

A more explicit expression for the cyclotomic polynomials is given by:

Lemma. *For each $n \geq 2$, we have*

$$\phi_n(T) = \prod_{\substack{1 \leq d < n \\ d|n}} v_{n/d}(T)^{\mu(d)}$$

4.3. Following [24] we say that a polynomial $p(T) \in \mathbb{Z}[T]$ is *represented* by $q(T) \in \mathbb{Z}[T]$ if

$$p(T^2) = q^*(T) := T^{\deg q} q(T + T^{-1}).$$

The interest in the representability of polynomials is due to the relation between the set of roots of $p(T)$ and $q(T)$ whenever $p(T^2) = q^*(T)$. Indeed, in that case $\text{Root } p(T) \subset \mathbb{S}^1$ (resp. $\mathbb{S}^1 \setminus \{1\}$) if and only if $\text{Root } q(T) \subset [-2, 2]$ (resp. $(-2, 2)$). In [24] is shown that the Coxeter polynomial $\hat{f}_{(p_1, \dots, p_t)}(T)$ of an extended canonical algebra of type $(p_1, \dots, p_t)$ is represented by

$$q_{(p_1, \dots, p_t)}(T) = T(T^2 - 1) \prod_{i=1}^t v_{p_i}(T) - \chi_{[p_1, \dots, p_t]}(T),$$

where $\chi_{[p_1, \dots, p_t]}(T)$ is the characteristic polynomial of the *adjacency matrix* of the star graph of type $[p_1, \dots, p_t]$.

Using the above expressions the following was recently shown by the authors:

Theorem [24]. *Let $A = C[P]$ be an extended canonical algebra of type $(p_1, \dots, p_{t-1}, p_t + 1)$ and A' be an extended canonical algebra of type $(p_1, \dots, p_t)$. Then the following holds:*

- (a) φ_A accepts at most 4 eigenvalues outside $\mathbb{S}^1$
- (b) If $\text{Root } f_A \subset \mathbb{S}^1$, then also $\text{Root } f_{A'} \subset \mathbb{S}^1$

Sketch of proof: The polynomials $q_{(p_1, \dots, p_t)}(T)$ satisfy a *Chebysheff recursion formula* as follows:

$$q_{(p_1, \dots, p_t+1)}(T) = Tq_{(p_1, \dots, p_t)}(T) - q_{(p_1, \dots, p_t-1)}(T).$$

A version of Sturm's Theorem applies to assure that for any real interval $[\alpha, \beta]$, if $q_{(p_1, \dots, p_t+1)}(T)$ has roots $\lambda_1 \leq \dots \leq \lambda_s$ in $[\alpha, \beta]$, then $q_{(p_1, \dots, p_t)}(T)$ has roots $\lambda'_1 \leq \dots \leq \lambda'_{s-1}$ in $[\alpha, \beta]$ satisfying

$$\lambda_1 \leq \lambda'_1 \leq \lambda_2 \leq \lambda'_2 \leq \dots \leq \lambda_{s-1} \leq \lambda'_{s-1} \leq \lambda_s.$$

(a) and (b) follow easily from these facts. □

4.4. According to (4.3), to prove Theorem 2 we need to calculate the minimal weight sequences p such that $\hat{f}_p(T)$ is not contained in $\mathbb{S}^1$. This is done by systematically computing the roots of Coxeter polynomials of extended canonical algebras.

Recall that the *spectral radius* of the Coxeter transformation φ_A is by definition $\rho_A(\varphi_A) = \max \{|\lambda| : \lambda \in \text{Root } f_A(T)\}$. In the following list A is an extended canonical algebra of weight type p . The invariant called *Dynkin index* is explained in section 5.

	Weight ($p_1, \dots, p_t$)	Irreducible factorization of $f_A(T)$	$\rho(\varphi_A)$	Dynkin index
$t=3$	(2, 3, 11)	$1+T-T^3-T^4+T^6+T^7+T^9+T^{10}-T^{12}-T^{13}+T^{15}+T^{16}$	1.1064	6
	(2, 4, 9)	$\phi_2\phi_5(T^{10}-T^9+T^5+T+1)$	1.1329	4
	(2, 5, 8)	$1+T+T^4+T^5+T^6+2T^8+T^9+T^{10}+T^{11}+T^{14}+T^{15}$	1.1574	4
	(2, 6, 7)	$1+T+T^4+2T^5+2T^6+T^7+T^8+2T^9+2T^{10}+T^{11}+T^{14}+T^{15}$	1.1669	4
	(3, 3, 8)	$1+T+T^2+T^5+2T^6+3T^7+2T^8+T^9+T^{12}+T^{13}+T^{14}$	1.1498	3
	(3, 4, 7)	$1+T+T^2+T^3+T^4+2T^5+3T^6+3T^7+3T^8+2T^9+T^{10}+T^{11}+T^{12}+T^{13}+T^{14}$	1.1847	3
	(3, 5, 6)	$\phi_3(T^{12}+T^9+T^8+T^7+T^6+T^5+T^4+T^3+1)$	1.1966	3
	(4, 4, 6)	$\phi_2\phi_4(T^{10}-T^9+T^8+T^6+T^4+T^2-T+1)$	1.2715	3
	(4, 5, 5)	$\phi_5(T^{10}+T^7+T^6+T^5+T^4+T^3+1)$	1.2277	3
$t=4$	(2, 2, 2, 7)	$\phi_2^2(T^{10}+T^6+T^5+T^4+1)$	1.1670	2
	(2, 2, 3, 6)	$\phi_2^2\phi_3(T^8-T^7+T^6+T^4+T^2-T+1)$	1.2196	2
	(2, 3, 3, 4)	$\phi_2\phi_4(T^8+T^6+T^5+2T^4+T^3+T^2+1)$	1.2874	2
	(3, 3, 3, 4)	$\phi_3^2(T^8+T^6+2T^5+2T^3+T^2+1)$	1.3307	2
$t=5$				2
	(2, 2, 2, 2, 5)	$\phi_2^3(T^8+T^6+T^3+2T^4+T^3+T^2+1)$	1.2874	2
	(2, 2, 2, 3, 4)	$\phi_2^3(T^8+2T^6+T^5+3T^4+2T^2+1)$	1.3351	2
	(2, 2, 3, 3, 3)	$\phi_2\phi_3^2(T^6+T^4+2T^3+T^2+1)$	1.3765	2
$t=6$				2
	(2, 2, 2, 2, 2, 3)	$\phi_2^4(T^6+2T^4+T^3+2T^2+1)$	1.3395	2
$t=7$				2
	(2, 2, 2, 2, 2, 2, 2)	$\phi_2^6(T^4-T^3+3T^2-T+1)$	1.5392	2

Table 1. Critical weight sequences.

4.5. To complete the proof of Theorem 2 we shall show that any weight sequence $p' < p$ with p in Table 1 has all its roots on $\mathbb{S}^1$. This is computed in the following Table 2.

As above $A = C[P]$ is an extended canonical algebra of weight type $p = (p_1, \dots, p_t)$. In Table 2, the marks $\bullet$ and $\square$ refer to the case $k = \mathbb{C}$: those weight sequences marked

by $\bullet$ or $\square$ correspond to algebras $R(p, \lambda)$ associated to hypersurface singularities, in those cases $R(p, \lambda)$ is formally 3-generated. The marks $\bullet$ correspond to *Arnold's 14 exceptional unimodal singularities* in the theory of singularities of differentiable maps [1]. Among those weight sequences $p = (p_1, p_2, p_3)$ (that is $t = 3$), Arnold's singularities are exactly those rings of automorphic forms having three generators [37].

Weight sequence	Factorization of $f_A(T)$	Poincaré series	Period of φ_A
• (2, 3, 7)	ϕ_{42}	(6, 14, 21) (42)	42
• (2, 3, 8)	$\phi_2 \cdot \phi_{10} \cdot \phi_{30}$	(6, 8, 15) (30)	30
• (2, 3, 9)	$\phi_3 \cdot \phi_{12} \cdot \phi_{24}$	(6, 8, 9) (24)	24
(2, 3, 10)	$\phi_2 \cdot \phi_{16} \cdot \phi_{18}$	(6, 8, 9, 10) (16, 18)	72
• (2, 4, 5)	$\phi_2 \cdot \phi_6 \cdot \phi_{30}$	(4, 10, 15) (30)	30
• (2, 4, 6)	$\phi_2^2 \cdot \phi_{22}$	(4, 6, 11) (22)	22
• (2, 4, 7)	$\phi_2 \cdot \phi_9 \cdot \phi_{18}$	(4, 6, 7) (18)	18
(2, 4, 8)	$\phi_2^2 \cdot \phi_4 \cdot \phi_{12} \cdot \phi_{14}$	(4, 6, 7, 8) (12, 14)	84
• (2, 5, 5)	$\phi_5 \cdot \phi_{20}$	(4, 5, 10) (20)	20
• (2, 5, 6)	$\phi_2 \cdot \phi_8 \cdot \phi_{16}$	(4, 5, 6) (16)	16
(2, 5, 7)	$\phi_{11} \cdot \phi_{12}$	(4, 5, 6, 7) (11, 12)	132
(2, 6, 6)	$\phi_2^2 \cdot \phi_3 \cdot \phi_6 \cdot \phi_{10} \cdot \phi_{12}$	(4, 5, 6, 6) (10, 12)	60
• (3, 3, 4)	$\phi_3 \cdot \phi_{24}$	(3, 8, 12) (24)	24
• (3, 3, 5)	$\phi_2 \cdot \phi_3 \cdot \phi_6 \cdot \phi_{18}$	(3, 5, 9) (18)	18
• (3, 3, 6)	$\phi_3^2 \cdot \phi_{15}$	(3, 5, 6) (15)	15
(3, 3, 7)	$\phi_2 \cdot \phi_3 \cdot \phi_4 \cdot \phi_{10} \cdot \phi_{12}$	(3, 5, 6, 7) (10, 12)	60
• (3, 4, 4)	$\phi_2 \cdot \phi_4 \cdot \phi_{16}$	(3, 4, 8) (16)	16
• (3, 4, 5)	ϕ_{13}	(3, 4, 5) (13)	13
(3, 4, 6)	$\phi_2 \cdot \phi_3 \cdot \phi_9 \cdot \phi_{10}$	(3, 4, 5, 6) (9, 10)	90
(3, 5, 5)	$\phi_2 \cdot \phi_5 \cdot \phi_8 \cdot \phi_{10}$	(3, 4, 5, 5) (8, 10)	40
• (4, 4, 4)	$\phi_2^2 \cdot \phi_4^2 \cdot \phi_6 \cdot \phi_{12}$	(3, 4, 4) (12)	12
(4, 4, 5)	$\phi_2 \cdot \phi_4 \cdot \phi_8 \cdot \phi_9$	(3, 4, 4, 5) (8, 9)	72
□ (2, 2, 2, 3)	$\phi_2^2 \cdot \phi_{18}$	(2, 6, 9) (18)	18
□ (2, 2, 2, 4)	$\phi_2^2 \cdot \phi_{14}$	(2, 4, 7) (14)	14
□ (2, 2, 2, 5)	$\phi_2^2 \cdot \phi_3 \cdot \phi_6 \cdot \phi_{12}$	(2, 4, 5) (12)	12
(2, 2, 2, 6)	$\phi_2^2 \cdot \phi_8 \cdot \phi_{10}$	(2, 4, 5, 6) (8, 10)	40
□ (2, 2, 3, 3)	$\phi_2 \cdot \phi_3 \cdot \phi_4 \cdot \phi_{12}$	(2, 3, 6) (12)	12
□ (2, 2, 3, 4)	$\phi_2^2 \cdot \phi_5 \cdot \phi_{10}$	(2, 3, 4) (10)	10
(2, 2, 3, 5)	$\phi_2 \cdot \phi_7 \cdot \phi_8$	(2, 3, 4, 5) (7, 8)	56
(2, 2, 4, 4)	$\phi_2^2 \cdot \phi_4 \cdot \phi_6 \cdot \phi_8$	(2, 3, 4, 4) (6, 8)	24
□ (2, 3, 3, 3)	$\phi_3^2 \cdot \phi_9$	(2, 3, 3) (9)	9
(2, 3, 3, 4)	$\phi_2 \cdot \phi_3 \cdot \phi_6 \cdot \phi_7$	(2, 3, 3, 4) (6, 7)	42
(3, 3, 3, 3)	$\phi_2 \cdot \phi_3^3 \cdot \phi_6^2$	(2, 3, 3, 3) (6, 6)	∞
□ (2, 2, 2, 2, 2)	$\phi_2^4 \cdot \phi_{10}$	(2, 2, 5) (10)	10
□ (2, 2, 2, 2, 3)	$\phi_2^3 \cdot \phi_4 \cdot \phi_8$	(2, 2, 3) (8)	8
(2, 2, 2, 2, 4)	$\phi_2^2 \cdot \phi_3 \cdot \phi_6^2$	(2, 2, 3, 4) (6, 6)	∞
(2, 2, 2, 3, 3)	$\phi_2^2 \cdot \phi_3 \cdot \phi_5 \cdot \phi_6$	(2, 2, 3, 3) (5, 6)	30
(2, 2, 2, 2, 2, 2)	$\phi_2^5 \cdot \phi_4 \cdot \phi_6$	(2, 2, 2, 3) (4, 6)	12

Table 2. Weights p with $\rho(\varphi_A) = 1$.

4.6. **Proof of Theorem 2.** (a) follows from (4.4) and (4.5) using Theorem (4.3). Part (b) is shown in [24], see (4.3). $\square$

5. The Poincaré series of an extended canonical algebra

5.1. Let $C = C(p, \lambda)$ be a wild canonical algebra with weight sequence $p = (p_1, \dots, p_t)$. Let P be an indecomposable projective C -module. We define the *Poincaré series* $\hat{P}_C = \hat{P}(p_1, \dots, p_t) \in \mathbb{Z}[[T]]$ by

$$\hat{P}_C(T) = \sum_{n=0}^{\infty} \langle [P], \varphi_C^n[P] \rangle_C T^n.$$

Recall that $\varphi_C^n[P] = [\tau_{D^b(\text{mod-}C)}^n P]$ in $K_0(D^b(\text{mod-}C)) = K_0(C)$. Moreover, observe that $T + \hat{P}_C(T)$ is the Hilbert-Poincaré series $P_C(T)$, as defined in the Introduction, for the graded algebra

$$R(p, \lambda) = \bigoplus_{n=0}^{\infty} \text{Hom}(L, \tau_{\mathbb{X}}^n L)$$

where $\mathbb{X} = \mathbb{X}(p, \lambda)$ is a weighted projective line, $\tau_{\mathbb{X}}$ is the Auslander-Reiten translation in $\text{coh}(\mathbb{X})$ and L is any rank one bundle, see [23] and [25]. In particular, $\hat{P}_C(T)$ does not depend on the choice of P .

Proposition. *With the notation above, let $A = C[P]$ be an extended canonical algebra. Then*

- (a) [25, Cor 3.6]: $f_A(T) = P_C(T)f_C(T)$
- (b) [23, Th. 8.6]: $P_C(T) = 1 + T - T \frac{f_{[p_1, \dots, p_t]}(T)}{f_{(p_1, \dots, p_t)}(T)}$
- (c) [23, Prop. 4.3]: $P_C(T) = T + \frac{1}{1-T} + (t-2) \frac{T}{(1-T)^2} - \sum_{i=1}^t \frac{T}{(1-T)(1-T^{p_i})}$. $\square$

5.2. We recall from [23] the following concepts.

Definition [23]. Assume $p = (p_1, \dots, p_t)$ is a weight sequence of wild type. The *Dynkin label* of p is the Dynkin diagram of one of the extended Dynkin graphs $[2, 2, 2, 2]$, $[3, 3, 3]$ [2, 4, 4] or $[2, 3, 6]$ specified as follows:

- (a) if $t \geq 4$, then the label is of type $[2, 2, 2, 2]$
- (b) if $t = 3$, then the label is of type $[a, b, c]$ if $[a, b, c] \leq [p_1, p_2, p_3]$ and $a + b + c$ is minimal.

We say that p has *Dynkin index* 2, 3, 4 or 6 if its Dynkin label is $[2, 2, 2, 2]$, $[3, 3, 3]$, $[2, 4, 4]$ or $[2, 3, 6]$ respectively.

Consider the graded algebra $R = R(p, \lambda)$. In [23] the *support monoid* $M(p)$ was introduced as the set of those $n \in \mathbb{N}$ with $R_n \neq 0$. Clearly, $M(p)$ is an additive semigroup in $\mathbb{N}$ generating $\mathbb{Z}$ as a group.

Proposition [23]. *The support monoid $M(p)$ is finitely generated with at most 6 generators. The smallest element in $M(p)$ is the Dynkin index of p .* $\square$

5.3. In a preliminary version of [23], the authors displayed the list of all possible support monoids $M(p)$. This list of 22 semigroups is essential for the proofs of Theorems 3 and 4 and we reproduce it below. For a weight sequence p the largest integer n such that n does not belong to $M(p)$ is called the *Frobenius number* of $M(p)$ and it is denoted by $\alpha(p)$. Of particular interest is the fact that $p \leq q$ implies $M(p) \subset M(q)$.

Weight type p	Frobenius number $\alpha(p)$	generators of $\mathbb{M}(p)$
(2, 3, 7)	43	$\{6, 14, 21\}$
(2, 3, 8)	25	$\{6, 8, 15\}$
(2, 3, 9)	19	$\{6, 8, 9\}$
(2, 3, 10)	13	$\{6, 8, 9, 10\}$
(2, 3, 11)	13	$\{6, 8, 9, 10, 11\}$
(2, 3, 12)	13	$\{6, 8, 9, 10, 11\}$
(2, 3, 13)	7	$\{6, 8, 9, 10, 11, 13\}$
(2, 3, ∞)	7	$\{6, 8, 9, 10, 11, 13\}$
(2, 4, 5)	21	$\{4, 10, 15\}$
(2, 4, 6)	13	$\{4, 6, 11\}$
(2, 4, 7)	9	$\{4, 6, 7\}$
(2, 4, 8)	9	$\{4, 6, 7\}$
(2, 4, 9)	5	$\{4, 6, 7, 9\}$
(2, 4, ∞)	5	$\{4, 6, 7, 9\}$
(2, 5, 5)	11	$\{4, 5\}$
(2, 5, 6)	7	$\{4, 5, 6\}$
(2, 5, 7)	3	$\{4, 5, 6, 7\}$
(2, 5, ∞)	3	$\{4, 5, 6, 7\}$
(2, 6, 6)	7	$\{4, 5, 6\}$
(2, 6, 7)	3	$\{4, 5, 6, 7\}$
(2, 6, ∞)	3	$\{4, 5, 6, 7\}$
(2, ∞ , ∞)	3	$\{4, 5, 6, 7\}$
(3, 3, 4)	13	$\{3, 8\}$
(3, 3, 5)	7	$\{3, 5\}$
(3, 3, 6)	7	$\{3, 5\}$
(3, 3, 7)	4	$\{3, 5, 7\}$
(3, 3, ∞)	3	$\{3, 5, 7\}$
(3, 3, 4)	5	$\{3, 4\}$
(3, 3, 5)	2	$\{3, 4, 5\}$
(4, 4, 4)	5	$\{3, 4\}$
(4, 4, 5)	2	$\{3, 4, 5\}$
(∞ , ∞ , ∞)	2	$\{3, 4, 5\}$
(2, 2, 2, 3)	7	$\{2, 9\}$
(2, 2, 2, 4)	5	$\{2, 7\}$
(2, 2, 2, 5)	3	$\{2, 5\}$
(2, 2, 2, ∞)	3	$\{2, 5\}$
(2, 2, 3, 3)	1	$\{2, 3\}$
(2, 2, 3, ∞)	1	$\{2, 3\}$
(∞ , ∞ , ∞ , ∞)	1	$\{2, 3\}$
(2, 2, 2, 2, 2)	3	$\{2, 5\}$
(2, 2, 2, 2, 3)	1	$\{2, 3\}$
(2, 2, 2, 2, ∞)	1	$\{2, 3\}$
(2, 2, 2, 2, 2, 2)	1	$\{2, 3\}$
(∞ , ∞ , $\dots$, ∞ , ∞)	1	$\{2, 3\}$

Table 3. Semigroups of $(\mathbb{N}, +)$ with the form $M(p)$.

5.4. For a given weight sequence $p = (p_1, \dots, p_t)$, the Coxeter polynomials $f_{(p_1, \dots, p_t)}(T)$ and $\hat{f}_{(p_1, \dots, p_t)}(T)$ are readily computed by (4.1). In case $\text{Root } \hat{f}_{(p_1, \dots, p_t)}(T) \subset \mathbb{S}^1$, then using (3.2), the Poincaré series $P_C(T)$ can be written as a rational function

$$P_C(T) = \frac{\prod_{i=1}^m (1 - T^{c_i})}{\prod_{j=1}^n (1 - T^{d_j})} (1 - T)^r$$

for sequences $(d_1, \dots, d_n)$ and $(c_1, \dots, c_m)$ of natural numbers ≥ 2 and some $r \in \mathbb{Z}$. In case p is of wild type, then by (4.1)

$$\hat{f}_{(p_1, \dots, p_t)}(1) = -f_{[p_1, \dots, p_t]}(1) = \left[(t-2) - \sum_{i=1}^t \frac{1}{p_i} \right] \prod_{i=1}^t p_i > 0$$

and by (5.1), $P_C(T)$ has a pole of order 2 at $T = 1$, that is $m - n + r = -2$. Moreover, developing the series $P_C(T)$ we readily see that $r \neq 0$ implies that the semigroups $M(p)$ is $\mathbb{N}$, but [23, Th. 10.4] claims that $1 + \left\lfloor \frac{1}{t-2} \right\rfloor \leq \alpha(p)$, that is, $\alpha(p) > 1$ and therefore $r = 0$. We state these considerations in the following.

Lemma. *Let $p = (p_1, \dots, p_t)$ be a weight sequence of wild type and $C = C(p, \lambda)$ be a canonical algebra. Then $\text{Root } \hat{f}_{(p_1, \dots, p_t)}(T) \subset \mathbb{S}^1$ if and only if $R(p, \lambda)$ is formally n -generated, that is*

$$P_C(T) = \frac{\prod_{i=1}^{n-2} (1 - T^{c_i})}{\prod_{j=1}^n (1 - T^{d_j})}$$

for numbers $c_1, \dots, c_{n-2}$ and $d_1, \dots, d_n$, all ≥ 2 , satisfying

$$1 + \sum_{j=1}^n d_j = \sum_{i=1}^{n-2} c_i.$$

Proof. If $\text{Root } \hat{f}_{(p_1, \dots, p_t)}(T) \subset \mathbb{S}^1$, we showed that $P_C(T)$ has the desired form. Moreover,

$$\sum_{j=1}^n d_j + \deg \hat{f}_{(p_1, \dots, p_t)}(T) = \sum_{i=1}^{n-2} c_i + \deg f_C(T).$$

The converse follows from $\hat{f}_{(p_1, \dots, p_t)}(T) = P_C(T)f_C(T)$. $\square$

In Table 2, for a weight sequence $p = (p_1, \dots, p_t)$ of wild type and $C = C(p, \lambda)$ the corresponding canonical algebra we have calculated (under the column ‘Poincaré series’) the sequences $(d_1, \dots, d_n)$ and $(c_1, \dots, c_{n-2})$ corresponding to $P_C(T) = \frac{\hat{f}_{(p_1, \dots, p_t)}(T)}{f_{(p_1, \dots, p_t)}(T)}$.

5.5. Proof of Theorem 3. Let $C = C(p, \lambda)$ be a canonical algebra of wild type and $A = C[P]$ be an extended canonical algebra. Implications (a) $\Rightarrow$ (b) $\Rightarrow$ (c) are clear.

(c) $\Rightarrow$ (a): Assume $\text{Root } f_A(T) \subset \mathbb{S}^1$. By (5.4), $P_C(T)$ is n -generated. Hence $M(p)$ is at most n -generated. By Table 3, if $n \geq 5$ then $p = (2, 3, p_3)$ with $p_3 \geq 11$. But a calculation shows (see Table 2) that $\text{Root } \hat{f}_{(2, 3, 11)}(T)$ is not contained in $\mathbb{S}^1$. Then

Theorem 2 implies that $\text{Root } \hat{f}_{(2,3,p_3)}(T)$ is not contained in $\mathbb{S}^1$ for $p_3 \geq 11$, which shows that $n \leq 4$.

For $k = \mathbb{C}$, the list entries in Table 2 correspond to Fuchsian singularities which are minimal elliptic as classified in [36]. These rings are graded complete intersection domains. $\square$

5.6. As a consequence of Theorem 3 and the classification given in Table 2 we get the following.

Corollary. *Let $C = C(p, \lambda)$ be a canonical algebra of wild type. Let $A = C[P]$ be an extended canonical algebra. The following are equivalent:*

- (a) φ_A is periodic.
- (b) $\text{Spec } \varphi_A \subset \mathbb{S}^1$ and p is not $(3, 3, 3, 3)$ or $(2, 2, 2, 2, 4)$.
- (c) $f_A(T) = \prod_{i=1}^m \phi_{s_i}(T)^{e_i}$ for $1 \leq s_1 < s_2 < \cdots < s_m$ and $e_i \geq 1$ ($1 \leq i \leq m$), with $e_m = 1$. $\square$

6. Graded integral domains with 3 homogeneous generators

6.1. The following simple remark is well-known.

Lemma. *Let R be a graded complete intersection integral k -algebra of Krull dimension 2. Then R is generated by 3 homogeneous elements if and only if $R = k[x_1, x_2, x_3]/(f)$ with $\deg x_i = d_i$ ($1 \leq i \leq 3$) and f a homogeneous prime polynomial. In this case the Poincaré-Hilbert series of R has the form*

$$\sum_{n=0}^{\infty} (\dim_k R_n) T^n = \frac{1-T^c}{(1-T^{d_1})(1-T^{d_2})(1-T^{d_3})}$$

for some natural numbers c, d_1, d_2, d_3 satisfying $1 + d_1 + d_2 + d_3 = c$.

Proof. Assume we have a graded surjection $g: k[x_1, x_2, x_3] \rightarrow R$ such that $y_i = g(x_i)$ is homogeneous of degree d_i , $1 \leq i \leq 3$. Since R is graded integral, then $I = \ker g$ is a prime ideal. Since R has Krull-dimension two, the ideal I has height one, hence it is principal. Let $I = (f)$ and $\deg(f) = c$. Then the Poincaré series has the desired form and $1 + d_1 + d_2 + d_3 = c$ by (4.4). $\square$

6.2. Proof of Theorem 4. Let $C = C(p, \lambda)$ be a wild canonical algebra and $A = C[P]$ a corresponding extended canonical algebra.

(a) $\Rightarrow$ (b): Assume $R(p, \lambda)$ is formally 3-generated. By (5.4), $\text{Root } f_A \subset \mathbb{S}^1$. The result follows from the list given in Table 2.

(b) $\Rightarrow$ (a): follows as above from Table 2 and (5.4).

Assume $R = R(p, \lambda)$ is formally 3-generated with

$$\sum_{n=0}^{\infty} (\dim_k R_n) T^n = \frac{1-T^c}{(1-T^{d_1})(1-T^{d_2})(1-T^{d_3})}$$

with $(d_1, d_2, d_3), (c)$ according to Table 2. We shall consider two distinguished situations:

if $t = 3$, then R is a quasi-homogeneous complete intersection of the form $k[x_1, x_2, x_3]/(f)$ with $\deg x_i = d_i$ and f a homogeneous relation as displayed in (6.3). The case $t \geq 4$ and $k = \mathbb{C}$ is considered in (6.4). $\square$

6.3. Theorem. [21]. *Let $C = C(p, \lambda)$ be a canonical algebra of wild weight type $p = (p_1, p_2, p_3)$ such that the graded algebra $R = R(p, \lambda)$ is formally 3-generated. Then R has the form*

$$R = k[x, y, z] = k[X, Y, Z]/(F)$$

where the relation F , the degree triple $\deg(x, y, z)$ and $\deg(F)$ are displayed in Table 4:

	p	$\deg(x, y, z)$	relation F	$\deg(F)$	Name
Index = 6	(2, 3, 7)	(6, 14, 21)	$Z^2 + Y^3 + X^7$	42	E_{12}
	(2, 3, 8)	(6, 8, 15)	$Z^2 + X^5 + XY^3$	30	Z_{11}
	(2, 3, 9)	(6, 8, 9)	$Z^2 + XZ^2 + X^4$	36	Q_{10}
4	(2, 4, 5)	(4, 10, 15)	$Z^2 + Y^3 + X^5Y$	30	E_{13}
	(2, 4, 6)	(4, 6, 11)	$Z^2 + X^4Y + ZY^3$	22	Z_{12}
	(2, 4, 7)	(4, 6, 7)	$Y^3 + X^3Y + XZ^2$	18	Q_{11}
	(2, 5, 5)	(4, 5, 10)	$Z^2 + Y^2Z + X^5$	20	W_{12}
	(2, 5, 6)	(4, 5, 6)	$XZ^2 + Y^2Z + X^4$	16	S_{11}
3	(3, 3, 4)	(3, 8, 12)	$Z^2 + Y^3 + X^4Z$	24	E_{14}
	(3, 3, 5)	(3, 5, 9)	$Z^2 + XY^3 + X^3Z$	18	Z_{13}
	(3, 3, 6)	(3, 5, 6)	$Y^3 + X^3Z + XZ^2$	15	Q_{12}
	(3, 4, 4)	(3, 4, 8)	$Z^2 - Y^2Z + X^4Y$	16	W_{13}
	(3, 4, 5)	(3, 4, 5)	$X^3Y + XZ^2 + Y^2Z$	13	S_{12}
	(4, 4, 4)	(3, 4, 4)	$X^4 - YZ^2 + Y^2Z$	12	U_{12}

$\square$

Table 4.

As observed in [21], these 14 equations are equivalent to Arnold's exceptional unimodal singularities. The equations are slightly different to those of the singularity theory classification, but equivalent for $k = \mathbb{C}$.

6.4. In view of the identification of $R(p, \lambda)$ with a ring of automorphic forms (5.5) in case $k = \mathbb{C}$, we get:

Theorem [9]. *Let $C = C(p, \lambda)$ be a canonical algebra of wild type $p = (p_1, \dots, p_t)$ with $t \geq 4$ over the complex numbers $\mathbb{C}$ and $R = R(p, \lambda)$ be the associated graded algebra. Then the following are equivalent:*

- (a) $R(p, \lambda)$ is formally 3-generated;
- (b) $9 \leq \sum_{i=1}^t p_i \leq 11$
- (c) *there is a parameter sequence $\lambda' = (\lambda'_3, \dots, \lambda'_t)$ such that the algebra $R(p, \lambda')$ is of the form*

$$k[x, y, z] = k[X, Y, Z]/(F)$$

where the relation F , the degree sequence $\deg(x, y, z)$ and $\deg(F)$ are displayed below:

	p	$\deg(x, y, z)$	relation F	$\deg(F)$	Name
$t = 4$	$(2, 2, 2, 3)$	$(2, 6, 9)$	$Z^2 + Y^3 + X^9$	18	$J_{3,0}$
	$(2, 2, 2, 4)$	$(2, 4, 7)$	$Z^2 + XY^3 + X^7$	14	$Z_{1,0}$
	$(2, 2, 2, 5)$	$(2, 4, 5)$	$Y^3 + XZ^2 + X^6$	12	$Q_{2,0}$
	$(2, 2, 3, 3)$	$(2, 3, 6)$	$Z^2 + Y^4 + X^6$	12	$W_{1,0}$
	$(2, 2, 3, 4)$	$(2, 3, 4)$	$Y^2Z + XZ^2 + X^5$	10	$S_{1,0}$
	$(2, 3, 3, 3)$	$(2, 3, 3)$	$Z^3 + Y^3 + X^3Y$	9	$U_{1,0}$
$t = 5$	$(2, 2, 2, 2, 2)$	$(2, 2, 5)$	$Z^2 + Y^5 + X^5$	10	$NA_{0,0}^1$
	$(2, 2, 2, 2, 3)$	$(2, 2, 3)$	$YZ^2 + Y^4 + X^4$	8	$VNA_{0,0}^1$

□

Table 5.

References

- [1] V.I. Arnold, S. Gusejn-Zade and A. Varchenko. *Singularities of differentiable maps*. Monographs in Math 82 Birkhäuser (1985).
- [2] M. Barot and H. Lenzing. *One-point extensions and derived equivalences*. J. Algebra 264 (2003) 1–5.
- [3] A.I. Bondal. *Representations of associative algebras and coherent sheaves*. Math. USSR-Izv. 34 (1990), 23–42.
- [4] A.I. Bondal and M.M. Kapranov. *Enhanced triangulated categories*. Math. USSR-Sb. 70 (1991), 93–107.

- [5] R. O. Buchweitz. *Maximal Cohen-Macaulay modules and Tate-cohomology over Gorenstein rings*. Unpublished manuscript (1987) 155 pp.
- [6] I. V. Dolgachev. *Automorphic forms and quasihomogeneous singularities*. Functional Anal. Appl. 9 (1975) 149–151.
- [7] W. Ebeling and C. T. C. Wall. *Kodaira singularities and an extension of Arnold’s strange duality*. Compositio Mathematica 56 (1985) 3–77.
- [8] W. Ebeling. The monodromy groups of isolated singularities of complete intersections. Lect. Notes Math. 1293, Springer, Berlin (1987).
- [9] W. Ebeling. *The Poincaré series of some special quasihomogeneous surface singularities*. Publ. RIMS, Kyoto Univ. 39 (2003), 393–413.
- [10] P. Gabriel. *Des catégories abéliennes*. Bull. Soc. Math. France 90 (1962), 323–448.
- [11] W. Geigle and H. Lenzing. *A class of weighted projective curves arising in representation theory of finite dimensional algebras*. In: Singularities, representation of algebras and vector bundles. Lect. Notes Math. 1273, Springer, Berlin (1987) 265–297.
- [12] W. Geigle and H. Lenzing. *Perpendicular categories with applications to representations and sheaves*. J. Alg. 144 (1991) 273–343.
- [13] S.I. Gelfand and Yu. I. Manin. *Methods of Homological Algebra*. Springer, Berlin (2003).
- [14] D. Happel. *Triangulated categories in the Representation Theory of finite dimensional algebras*. London Math. Soc. Lect. Note Ser. 119 (1988).
- [15] D. Happel, I. Reiten, S. Smalø, *Tilting in abelian categories and quasitilted algebras*, Mem. Amer. Math. Soc. 575 (1996).
- [16] D. Happel. *A characterization of hereditary categories with tilting object*. Invent. Math. 144 (2001) 161–179.
- [17] H. Kajiura, K. Saito and A. Takahashi. *Matrix factorizations and representations of quivers II: Type ADE case*. arXiv:math.AG/0511155v2 3 Apr 2006.
- [18] B. Keller. *On differential graded categories*. Preprint 2006.
- [19] H. Krause. *The stable derived category of a noetherian scheme*. Compositio Math. 141 (2005), 1128–1162.
- [20] H. Laufer. *On minimally elliptic singularities*. Ann. J. of Math. 99 (1977) 1257–1295.
- [21] H. Lenzing. *Wild canonical algebras and rings of automorphic forms*. In: Finite dimensional algebras and related topics. Kluwer (1994) 191–212.
- [22] H. Lenzing and H. Meltzer. *Tilting sheaves and concealed-canonical algebras*. In: Proceedings of ICRA VII. CMS Conf. Proc. 18 (1996) 455–473.
- [23] H. Lenzing and J. A. de la Peña. *Wild canonical algebras*. Math. Z 224 (1997) 403–425.
- [24] H. Lenzing and J. A. de la Peña. *Coxeter polynomials and a Chebysheff recursion formula*. To appear.
- [25] H. Lenzing. *Coxeter transformations associated with finite dimensional algebras*. In: Progress in Math 173 (1999) 287–308.
- [26] J. Milnor. *On the 3 dimensional Brieskorn manifolds $m(p, q, r)$* . In: Knots, groups and 3-manifolds; Pap. dedic. Mem. R.H. Fox (1975) 175–225.
- [27] A. Neeman. *Triangulated Categories*. Annals of Mathematics Studies, 148. Princeton University Press, Princeton (2001).
- [28] W. Neumann. *Brieskorn complete intersections and automorphic forms*. Invent. Math. 42 (1977) 285–293.
- [29] D. Orlov. *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*. Proc. Steklov Inst. Math. 246 (2004) 227–248.
- [30] D. Orlov. *Derived categories of coherent sheaves and triangulated categories of singularities*. arXiv:math.AG/0503632v2 22 Sep 2005.

- [31] J. A. de la Peña. *On the dimension of the module-varieties of tame and wild algebras*. Comm. in Algebra 19 (6), (1991) 1795–1807.
- [32] J. Rickard. *Morita theory for derived categories*. J. London Math. Soc. 39 (1989), 436–456.
- [33] C. M. Ringel. *The canonical algebras*. In: Topics in Algebra, part I. Banach Center Pub. 26 (1950) 407–432.
- [34] C. M. Ringel. *Tame algebras and integral quadratic forms*. Lect. Notes Math., Springer, Berlin 1099 (1984).
- [35] K. Ueda. *Homological mirror symmetry and simple elliptic singularities*. arXiv:math.AG/0604361 v2 3 Jul 2006.
- [36] P. Wagreich. *Automorphic forms and singularities with C^* -action*. Illinois J. Math. 25 (1981) 359–382.
- [37] P. Wagreich. *Algebras of automorphic forms with few generators*. Trans. Amer. Math. Soc. 262 (1980) 367–389.

H. Lenzing
Institut für Mathematik
Universität Paderborn
33095 Paderborn
Germany
helmut@math.uni-paderborn.de

J. A. de la Peña
Instituto de Matemáticas
Universidad Nacional Autónoma de México
Ciudad Universitaria
México 04510, D. F.
México
jap@matem.unam.mx